arXiv:hep-ph/0109015v1 3 Sep 2001

Lepton flavor violating process in a supersymmetric grand unified theory with right-handed neutrino [⋆]

Seungwon Baek [a] Toru Goto [b] Yasuhiro Okada [b,c]
Ken-ichi Okumura [d]

[a] *Department of Physics, National Taiwan University, Taipei 106, Taiwan*

[b] *Theory Group, KEK, Tsukuba, Ibaraki, 305-0801, Japan*

[c] *Department of Particle and Nuclear Physics, The Graduate University of Advanced Studies, Tsukuba, Ibaraki, 305-0801, Japan*

[d] *Institute for Cosmic Ray Research, University of Tokyo, Kashiwanoha 5-1-5, Kashiwa, 277-8582, Japan*

Abstract

Motivated from the recent results of neutrino oscillation experiment, we investigated lepton flavor violating (LFV) processes in a SU(5) supersymmetric grand unified theory with right-handed neutrino. The current experimental upper bound for $\mu \to e\gamma$ process gives already some constraint on the model. Correlation between $\mu \to e\gamma$ and the SUSY contribution to the muon anomalous magnetic moment is also discussed. Future LFV experiments will give considerable impacts on this type of SUSY GUT models equipped with seesaw neutrino mass generation.

Key words: lepton flavor violation, SUSY GUT, neutrino oscillation

---

Supersymmetric standard model (SUSY SM) is a promising candidate for physics beyond the SM. Supersymmetry resolves elegantly the hierarchy problem of the SM. Furthermore, gauge coupling unification is realized if we assume grand unification with the SU(5) gauge group or its extensions. On the other hand, the seesaw mechanism naturally explain the smallness of neutrino mass which is suggested from the recent neutrino oscillation experiments. It introduces heavy Majorana right-handed neutrinos and their Yukawa interaction to lepton doublets. In the SUSY models, it is known that a radiative correction to the slepton mass matrix from this Yukawa inter-

---

[⋆] Talk given by K. Okumura at the 3rd International Workshop on Neutrino Factories based on Muon Storage Rings (NuFACT'01 Workshop) Tsukuba, Japan, May 24-30, 2001 action induces lepton flavor violating (LFV) processes such as $\mu \to e\gamma$ which is forbidden in the SM.

In this paper, we show the results of our numerical analysis for the LFV processes in a SU(5) supersymmetric grand unified theory (SUSY GUT) with right-handed neutrino [1]. We calculated branching ratios of $\mu \to e\gamma$, $\tau \to \mu\gamma$, and $\tau \to e\gamma$, and the SUSY contribution to the muon anomalous magnetic moment, for which recently BNL E821 experiment reported 2.6σ deviation from the SM theoretical calculation (2). In our analysis, we introduced higher dimensional operators above the GUT scale to accommodate realistic mass relations between the down-type quarks and the charged leptons. Consequently, new degrees of freedom appear at the GUT scale when we embed the fields of minimal SUSY SM (MSSM) in the SU(5) multiplets as $\mathbf{10}^i[Q^i, (V_U)^i_j\overline{U}^j, (V_E)^i_j\overline{E}^j]$, $\overline{\mathbf{5}}^i[(V_D)^i_j\overline{D}^j, L^i]$ where V_U, V_E, and V_D are unitary matrices defined in the basis where the mass matrices of down-type quark and charged lepton are diagonal. In the actual analysis, we considered the effects of two mixing angles θ_E in V_E and θ_D in V_D which describe the rotation between the first and the second generations. We numerically solved renormalization group equations from the Planck scale to the EW scale. For the initial condition, we assumed universal scalar and gaugino masses m_0, M_0 and SUSY breaking scalar couplings are taken to be proportional to the superpotential with a universal coefficient $m_0 A_0$.



Fig. 1. Contour plots of a_μ^{SUSY} and $B(\mu \to e\gamma)$ in the m_0-M_0 plane.

Fig.1 (a) and (b) shows the contour plots of the SUSY contribution to the muon anomalous magnetic moment a_μ^{SUSY} and $B(\mu \to e\gamma)$ in the m_0-M_0 plane. We take $\tan\beta = 20$, $\mu > 0$, $A_0 = 0$, $\theta_D = \theta_E = 0$, and $M_R = 4.0 \times 10^{13}$ GeV where M_R is the Majorana mass scale of the right-handed neutrinos. We assumed the MSW large mixing angle solution for the solar neutrino anomaly. The shaded region is excluded by the direct SUSY searches at LEP and Tevatron, the Higgs boson search at LEP, the result of $B(b \to s\gamma)$ at CLEO, and $B(\mu \to e\gamma)$. The current experimental upper bound, $B(\mu \to e\gamma) < 1.2 \times 10^{-11}$ already gives some constraint on the model. The favored region of BNL E821 experiment which corresponds to $20 \times 10^{-10} \lesssim a_\mu^{SUSY} \lesssim 60 \times 10^{-10}$ still survives within these constraints. An experimental improvement by order of magnitude could change the situation. If we lower M_R, the $\mu \to e\gamma$ constraint becomes weak while a_μ^{SUSY} is intact, however, in such a case, $y_{\nu 3}$ becomes much smaller than y_t and further GUT unification scenario such as SO(10) SUSY GUT becomes difficult. If we lower $\tan\beta$, not only $B(\mu \to e\gamma)$ but also a_μ^{SUSY} decreases and the constraint from the Higgs boson search becomes stronger. With the same parameter set, $B(\tau \to \mu\gamma)$ is $\lesssim 10^{-8}$ and $B(\tau \to e\gamma)$ is far below the current experimental bound. μ-e conversion in atomic nuclei is an another important LFV process. When the process is dominated by the photon-penguin diagrams as in our case, it is known that the conversion rate is related to $\mu \to e\gamma$ as $B(\mu^- N \to e^- N)/B(\mu^+ \to e^+\gamma) \approx B(A,Z)/428$ where $B(A,Z) \approx 1.1$ for ^{27}Al, $B(A,Z) \approx 1.8$ for ^{48}Ti and $B(A,Z) \approx 1.25$ for ^{208}Pb.

---

[1] This talk is based on the work in Ref.(1).

Fig.1 (c) and (d) shows the same plots for the case $\theta_D = \pi/4$. Other parameters are same as (a) and (b). Similar plots can be drawn for $\theta_E = \pi/4$. Inclusion of the effects of higher dimensional operators generally increases $B(\mu \to e\gamma)$ especially in the small m_0 region. The parameter region where the BNL E821 experiment favors is excluded in this figure. It is found that this contribution does not disappear even if we lower M_R because it mainly comes from the flavor mixing in LR mixing mass which originates in the matching condition of SUSY breaking scalar couplings at the GUT scale rather than the radiative corrections from the neutrino Yukawa coupling. $B(\tau \to \mu\gamma)$ is not affected by θ_D and θ_E and $B(\tau \to e\gamma)$ remains negligible. Fig.2 shows a correlation between the a_μ^{SUSY} and the $B(\mu \to e\gamma)$ where we searched SUSY parameters as $m_0, M_0 < 3$ TeV, $|A_0| < 5$. We also took both sign of μ parameter. Other parameters are same as Fig.1 (a) and (b). The dotted lines show the deviation reported by the BNL E821 experiment and the vertical line indicates the current experimental upper bound for $B(\mu \to e\gamma)$. It is clearly seen that future LFV experiments will give considerable impacts on this type of SUSY GUT models combined with the further improvement of our understanding on the muon anomalous magnetic moment.



Fig. 2. Correlation between a_μ^{SUSY} and $B(\mu \to e\gamma)$.

In summary, LFV processes provide us a unique opportunity testing the physics at the GUT scale and the Majorana mass scale which is relevant with the origin of neutrino masses and mixing and future experimental progress is expected to bring some insights on it.

References

[1] S. Baek, T. Goto, Y. Okada, and K. Okumura, Phys. Rev. D**63**,

051701 (2001); hep-ph/0104146, to appear in PRD, (See also references therein).

[2] Muon g-2 Collaboration, H.N. Brown *et al.*, Phys. Rev. Lett. **86**, 2227 (2001).